



Why formation rate of 3 nm particles depends linearly on sulphuric acid concentration?

M. Kulmala, K. E. J. Lehtinen, A. Laaksonen

► To cite this version:

M. Kulmala, K. E. J. Lehtinen, A. Laaksonen. Why formation rate of 3 nm particles depends linearly on sulphuric acid concentration?. *Atmospheric Chemistry and Physics Discussions*, 2005, 5 (6), pp.11277-11293. hal-00301906

HAL Id: hal-00301906

<https://hal.science/hal-00301906>

Submitted on 8 Nov 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Formation rate of
3 nm particles**

M. Kulmala et al.

Why formation rate of 3 nm particles depends linearly on sulphuric acid concentration?

M. Kulmala¹, K. E. J. Lehtinen², and A. Laaksonen³

¹University of Helsinki, Department of Physical Sciences, P.O. Box 64, FIN-00014 Helsinki, Finland

²Finnish Meteorological Institute and University of Kuopio, Department of Applied Physics, P.O. Box 1627, FIN-70211 Kuopio, Finland

³University of Kuopio, Department of Applied Physics, P.O. Box 1627, FIN-70211 Kuopio, Finland

Received: 7 October 2005 – Accepted: 29 October 2005 – Published: 8 November 2005

Correspondence to: M. Kulmala (markku.kulmala@helsinki.fi)

© 2005 Author(s). This work is licensed under a Creative Commons License.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

Abstract

According to atmospheric observations new particle formation seems to be a function of sulphuric acid concentration to the power from one to two. The nucleation theorem then predicts that the critical cluster contains one to two sulphuric acid molecules.

5 However, existing nucleation theories predicts that the power is more (or equal) than 2. Here we present an activation theory, which can explain the observed slope. In cluster activation the clusters containing one sulphuric acid molecule will activate for further growth due to heterogeneous nucleation, heterogeneous chemical reactions including polymerization or activation of soluble clusters. In the activation process
10 organic vapours are typically needed as condensing agents.

1. Introduction

Aerosol particles are ubiquitous in the Earth's atmosphere and affect our quality of life in many different ways. In polluted urban environments, aerosol emissions can affect human health through their inhalation (e.g. Donaldson et al., 1998), whilst globally,
15 aerosols are thought to contribute to climate change patterns (e.g. Charlson et al., 1987; Lohmann and Feichter, 2005). In recent years, considerable effort has been devoted to understanding how aerosols directly affect the Earth's radiation budget by scattering and absorbing incoming solar radiation. Aerosols also affect the radiation budget indirectly by modifying many cloud properties such as their albedo and lifetime.
20 It is generally thought that increases in aerosol concentrations will lead to brighter and more sustained clouds, thus providing additional planetary cooling.

In order to be able to better understand the health and climatic effects of atmospheric aerosols, the formation and growth processes of atmospheric aerosols should also be better understood (Kulmala, 2003). Nucleation, the formation of ultrafine particles detected at a few nm, and subsequent growth to ~100 nm in 1–2 days, has been
25 observed frequently in the continental boundary layer. Such observations span from

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Formation rate of
3 nm particles**

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

northern-most sub-arctic Lapland (Vehkamäki et al., 2004), over the remote boreal forest (Mäkelä et al., 1997; Kulmala et al., 1998, 2001a) and suburban Helsinki (Väkevää et al., 2000), to industrialised agricultural regions in Germany (Birmili and Wiedensohler, 1998) and also to coastal environments around Europe (O'Dowd et al., 1999).

The atmospheric new particle formation rates have also been investigated by Weber et al. (1996, 1997), and the biogenic aerosol formation by Kavouras et al. (1998). A recent overview summarised the formation and growth properties in a global point of view (Kulmala et al., 2004a), quantifying especially the formation and growth rates of nucleation events where available.

It has been proposed and also observed that atmospheric new particle formation depends on the sulphuric acid concentration. In laboratory experiments this dependence is clear and straightforward (Viisanen et al., 1997; Bernd et al., 2005), and the dependence is strong, in power-law form having exponents of order 5–10. In atmospheric conditions the dependence is much smoother. In the present paper we investigate the theoretical background of the observed sulphuric acid concentration dependence. For this purpose, we derive a theory for cluster activation, and compare its predictions with kinetic nucleation and observed particle formation.

2. Relations between fresh particle formation rate and sulphuric acid concentration

Critical clusters – if electrically neutral – formed by atmospheric nucleation events cannot yet be measured quantitatively due to instrumental limitations. Only one measurement of clusters during nucleation events has been reported, and it showed that clusters were present when 2.7–4 nm particles were detected (Weber et al., 1995). More work on the distribution and composition of such clusters is needed to refine our understanding of atmospheric nucleation. However, ion clusters can be and have been measured during nucleation events (e.g. Hörrak, 2001; Laakso et al., 2004)

Because critical neutral clusters cannot yet be measured, we are unable to measure

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

the true atmospheric nucleation rate but rather the formation rate of particles of some larger diameter d_p . The diameter d_p corresponds typically to the CNC detection limit, which is presently 3 nm or greater. In any case, based on the observed formation rates, the nucleation rate (formation rate of 1 nm particles) can be estimated based on theory describing the competition between condensation growth and cluster scavenging (Kerminen and Kulmala, 2002).

Using the observed nucleation mode growth rate and condensation sink determined from size spectra measurements, the concentration of condensable vapour and its source rate during the nucleation and growth events can be analysed (see Kulmala et al., 2001b). The aerosol condensation sink determines how rapidly molecules will condense onto pre-existing aerosols and depends strongly on the shape of the size distribution (see e.g. Pirjola et al., 1999; Kulmala et al., 2001b). The condensation sink CS is obtained by integrating over the aerosol size distribution:

$$CS = 2\pi D \int_0^{\infty} d_p \cdot \beta_M(d_p) \cdot n(d_p) d d_p = 2\pi D \sum_i \beta_{M_i} d_{p,i} N_i, \quad (1)$$

where $n(d_p)$ is the particle size distribution function and N_i is the concentration of particles in the size section i . Here d_p is particle diameter, D is diffusion coefficient of condensable vapour, and for the transitional correction factor for the mass flux β_m we use the Fuchs-Sutugin expression (Fuchs and Sutugin, 1971).

The measured formation rate is smaller than the actual nucleation rate J^* and determined by the competition between condensational growth (GR = growth rate from experimental observations) and scavenging (rate proportional to condensation sink CS) (Kerminen and Kulmala, 2002):

$$J_m = J^* \exp \left\{ 0.23 \left(1/d_m - 1/d^* \right) \frac{CS}{GR} \right\} \quad (2)$$

Setting the formation at 3 nm (J_3) and nucleation at 1 nm (J_1) gives

$$J_3 = J_1 \exp \left\{ -0.153 \frac{CS}{GR} \right\} \quad (3)$$

It is important to note that in deriving Eq. (2) (and thus also 3) a constant growth rate GR has been assumed. It has been seen, however, that the growth rate in the range 1–3 nm may be size dependent, caused presumably by organic vapour Kelvin effects (Kulmala et al., 2004b; Hirsikko et al., 2005). Since, in addition, Eq. (3) may be quite sensitive on GR , it should be considered as order-of-magnitude estimate only.

It has been seen (e.g. Weber et al., 1996) that the formation rate of 3 nm particles can be expressed as a power-law dependence of the sulphuric acid concentration

$$J_3 \propto [\text{H}_2\text{SO}_4]^{n_3} \quad (4)$$

with a power-law exponent n_3 . Correspondingly, we can assume that the formation rate of 1 nm particles has a similar dependence:

$$J_1 \propto [\text{H}_2\text{SO}_4]^{n_1} \quad (5)$$

By taking logarithms on each side, Eq. (3) can be rewritten as

$$\log(J_3) = \log(J_1) - 0.153 \frac{CS}{GR} = n_1 \log[\text{H}_2\text{SO}_4] - 0.153 \frac{CS}{GR} \quad (6)$$

Now, in addition to the nucleation rate terms J_1 and J_3 , the sulphuric acid concentration $[\text{H}_2\text{SO}_4]$ can appear in Eq. (6) in the growth rate term GR . The formation rate dependency on $[\text{H}_2\text{SO}_4]$ is illustrated in Fig. 1. The curve representing J_1 is obviously a straight line, with the slope n_1 depending on the nucleation mechanism. If the effect of sulphuric acid on growth is negligible, i.e., the growth rate GR does not depend on $[\text{H}_2\text{SO}_4]$, then the term $0.153 \frac{CS}{GR}$ in Eq. (8) is constant (with respect to $[\text{H}_2\text{SO}_4]$). Hence the resulting apparent nucleation rate at 3 nm J_3 (Fig. 1) has the same slope as J_1 , but a lower absolute value than J_1 by magnitude $0.153 \frac{CS}{GR}$. If GR depends on $[\text{H}_2\text{SO}_4]$, the exact value for the slope is not straightforward to determine. In any

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

case, it is clear that at otherwise the same conditions, increasing $[\text{H}_2\text{SO}_4]$ will increase GR , thus decreasing the magnitude of $0.153\text{ CS}/GR$ (see also Laaksonen, 2000). This means that J_3 will approach J_1 at high values of $[\text{H}_2\text{SO}_4]$, as indicated qualitatively (dashed line) in Fig. 1. Thus the slope for J_3 should be steeper than the slope for J_1 , i.e.,

$$n_1 \leq n_3. \quad (7)$$

The atmospheric measurements made by Weber et al. (1996) show that the slope for J_3 was between 1 and 2 in Idaho Hill. The recent measurement campaign (QUEST 2) (see Kulmala et al., 2004b) in Hyytiälä, Finland showed also that the slope is smaller than 2. In Fig. 2 the sulphuric acid concentration and the concentration of particles in size range 3–6 nm are plotted for one representative day of the campaign. The sulphuric acid concentration is scaled and also shifted 90 min ahead in time. This indicates the time needed for growth from nucleated size to 3 nm by condensation. The patterns of the curves are remarkably similar, indicating a linear relationship (i.e. a power-law exponent of unity) between these variables. Not all event days during the campaign show such a clear linear dependence. We can, however, confidently state that in all cases the power-law exponent is between one and two.

According to the nucleation theorem the slope determines the number of molecules in the critical cluster (Kashchiev, 1982):

$$[\partial \ln(J)/\partial \ln(A_i)]_{T,A_j} = n_i + \Delta, \quad (8)$$

where J denotes nucleation rate, A_i is the gas-phase activity (partial pressure divided by saturation vapor pressure) of species i , n_i is the number of molecules in the critical cluster, and Δ is a small term between 0 and 1 (Oxtoby and Kashchiev, 1994). It has been shown using both thermodynamic (Oxtoby and Kashchiev, 1994) and statistical mechanical (Viisanen et al., 1993; MacDowell, 2003) arguments that the nucleation theorem is a very general relation that extends down to the smallest cluster sizes and holds independently of any specific nucleation theories. Note that the derivatives

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

should be taken at constant temperature and gas phase activities of other species participating in the nucleation process, and in atmospheric conditions, also at constant condensation sink.

Based on our observations the number n_1 must be pretty small, approaching unity. It is smaller than n_3 , and n_3 is already two or smaller. This brings us to the question what is the mechanism responsible for n_1 being (at least near to) unity? Kinetic (barrier less) nucleation predicts that n_1 is 2, thermodynamic nucleation like binary (water + sulphuric acid) predicts values bigger than 10, and ternary nucleation values between 5 and 10 (see e.g. Bernd et al., 2005). How is it thus possible that $n_3=1-2$, as observed in the atmosphere, particularly in the continental boundary layer, although n_3 should be bigger than n_1 , and according to above mentioned theories n_1 is more or equal than 2. One possible explanation is multicomponent nucleation with many more compounds participating than 3, and the other one is activation of existing clusters.

3. Activation of clusters

In the atmosphere ion clusters are practically always and everywhere present (see e.g. Laakso et al., 2004). Some studies predict also the existence of neutral clusters (e.g. Kulmala et al., 2000; Kulmala et al., 2005). Therefore, in principle, activation on neutral or ion clusters can explain the observed values of n_3 .

The simplest way to describe the activation process is by using the concept of activation probability, such as in the theory of heterogeneous nucleation (see e.g. Lazaridis et al., 1992).

The time evolution of an activating cluster concentration can be estimated as

$$\frac{dN_{\text{clusters}}}{dt} = -kN_{\text{clusters}} \quad (9)$$

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

resulting in the well-known expression for activation probability P :

$$P = \frac{N_{\text{activated}}}{N_{\text{clusters}}} = 1 - \exp(-kt). \quad (10)$$

Here $N_{\text{activated}}$ is the number of activated clusters or actually aerosol particles, and N_{cluster} is the number concentration of clusters. k is a proportionality coefficient which is related to either a) heterogeneous nucleation, b) Köhler type activation or c) heterogeneous chemical reactions including polymerization.

In other words, Eq. (9) means that the activation rate J_{act} is:

$$J_{\text{act}} = kN_{\text{clusters}}$$

And the formation rate at 3 nm resulting from the activation mechanism can be estimated similarly as in Eq. (2) as:

$$J_3 = J_{\text{act}} \exp \left\{ 0.23 \left(1/3 - 1/d_{\text{act}} \right) \frac{CS}{GR} \right\} = kN_{\text{clusters}} \exp \left\{ 0.23 \left(1/3 - 1/d_{\text{act}} \right) \frac{CS}{GR} \right\} \quad (11)$$

In which d_{act} is the size at which activation is occurring.

Now it is evident that if ion or neutral clusters containing 1 sulphuric acid molecule will activate the observed slope of unity is explained. As already mentioned ion clusters are ubiquitous. In addition, large amounts of neutral ammoniumbisulphate clusters have been theoretically recently predicted (Vehkamäki et al., 2004).

In the following we investigate how different theories used in nucleation will work as compared with observed number concentrations.

In the test we compare 3 different ways to estimate formation rates and compared it with the formation rate calculated from observed number concentration of 3–6 nm particles taking into account losses due to coagulation and condensation growth out of size range.

$$J_{3,1} = C_1 [\text{H}_2\text{SO}_4] \exp \left\{ 0.23 \left(1/3 - 1 \right) \frac{CS}{GR} \right\} \quad (12a)$$

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

$$J_{3,2} = C_2 [\text{H}_2\text{SO}_4]^2 \exp \left\{ 0.23 \left(\frac{1}{3} - 1 \right) \frac{CS}{GR} \right\} \quad (12b)$$

$$J_{3,3} = C_3 [\text{H}_2\text{SO}_4]^3 \exp \left\{ 0.23 \left(\frac{1}{3} - 1 \right) \frac{CS}{GR} \right\} \quad (12c)$$

Equation (12a) describes activation theory, (12b) kinetic nucleation and (12c) is a conservative slope estimate for thermodynamic theories in general (slope value 3, which is smaller than typical value for thermodynamic nucleation).

The results are presented in Fig. 3. The best agreement seems to be with activation theory, i.e. Eq. (12a). The thermodynamic theory (Eq. 12c) gives clearly the worst predictions. From the fitted pre-factor C_i values one can also obtain estimates for the activation coefficient k ($=C_1$) and also for the kinetic pre factor (C_2), if kinetic nucleation is assumed. The typical growth time is 1000–10 000 s, and the GR for the day shown is 1.4 nm/h. The obtained values for the pre factors C_i are $C_1 \sim 5e-7$ (Activation). $C_2 \sim 2e-13$ and $C_3 \sim 6e-20$. As a general result we can say that the present activation theory explains why we can see from atmospheric observations slope of unity. However, as seen from figure, all theories underestimate measured formation rates before noon. This means that e.g. activation coefficient should be a factor of 10–100 higher than given at that time. Taking into account observed overcharging during that morning (see Laakso et al., 2004), this indicates that activation coefficient for ion activation is higher and ions are activated first (before noon) and later also neutral clusters.

Quite often organic vapours are needed for activation. The organic vapour can participate in the activation processes via a) heterogeneous reactions (Claeys et al., 2004), b) polymerization (Kalberer et al., 2004), c) heterogeneous nucleation or Nano-Köhler (see Kulmala et al., 2004c). Actually, the observed k value for the activation mechanism can be explained by product of concentration of collided molecules (around $1e7-1e8 \text{ cm}^{-3}$) and typical chemical reaction rate in gas phase.

As mentioned above, in the activation process, there are other vapours present as sulphuric but what is their contribution to the observed aerosol formation rate J_3 ? In

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

the case of multicomponent homogeneous nucleation, the nucleation theorem will tell us, what would be their contribution. In the case of heterogeneous nucleation the contribution will depend on contact angle and the ratio of sizes of seed particle and critical cluster (Määttä et al., 2005; Kulmala et al., 2001; Vehkamäki et al., 2005).

5 The values are often ten times smaller than in homogeneous case.

4. Conclusions

Current estimates on the magnitude of regional and global secondary aerosol formation rely almost entirely on modeling and are subject to large uncertainties. There are uncertainties in identifying both the detailed nucleation mechanisms as well as the nucleation rates. In this paper we present a method with which we can obtain hints about the formation mechanisms indirectly: by comparing sulfuric acid concentrations with particle concentrations between 3 nm and 6 nm, clear similarities are observed in their time evolution. The particle concentration seems to have a power-law dependence on the sulfuric acid concentration, with an exponent value between 1 and 2. A theoretical investigation of the connection between different exponent values and nucleation mechanisms shows that particle nucleation proceeds likely by either a kinetic mechanism or by activation mechanism.

The activation mechanism is based on the concept of activation probability. This is commonly used in studies of heterogeneous nucleation (see e.g. Kulmala et al., 2001; Wagner et al., 2003). In practice the activation probability can depend on, in addition to heterogeneous nucleation, also on heterogeneous chemical reactions including polymerization and activation of mutual soluble compounds, like in nano-Köhler theory. The activation constant is in the order of 10^{-6} – 10^{-7} 1/s. In the proposed activation mechanism neutral or ion clusters containing one sulphuric acid molecule are activated for further growth. This can directly explain the observed relationship between sulphuric acid concentration and fresh particle formation with slope unity.

Acknowledgements. Discussions and help by V.-M. Kerminen and S.-L. Sihto and the financial

11286

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

support from the Academy of Finland are acknowledged.

References

- Bernd, T., Böge, O., Stratmann, F., Heintzenberg, J., and Kulmala, M.: Rapid formation of new sulfuric acid particles at near-atmospheric conditions, *Science*, 307 (5710), 698–700, 2005.
- 5 Birmili, W. and Wiedensohler, A.: The influence of meteorological parameters on ultrafine particle production at a continental site, *J. Aerosol Sci.*, 29, S1015–1016, 1998.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655–661, 1987.
- 10 Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M., Artaxo, P., and Maenhaut, W.: Formation of secondary organic aerosols through photooxidation of isoprene, *Science*, 303, 1173–1176, 2004.
- Donaldson, K., Li, X. Y., and MacNee, W.: Ultrafine (nanometre) particle mediated lung injury, *J. Aerosol. Sci.*, 29(5-6), 553–560, 1998.
- Fuchs, N. A. and Sutugin, A. G.: Highly dispersed aerosol, in: *Topics in Current Aerosol Research*, edited by: Hidy, G. M. and Brock, J. R., Pergamon, New York, 1971.
- 15 Hirsikko, A., Laakso, L., Horrak, U., Aalto, P. P., Kerminen, V.-M., and Kulmala, M.: Annual and size dependent variation of growth rates and ion concentrations in boreal forest, *Boreal Env. Res.*, 10, 357–369, 2005.
- Hörrak, U.: *Air Ion Mobility Spectra at a rural area*, PhD thesis, University of Tartu, Estonia, 2001.
- 20 Kavouras, I. G., Mihalopoulos, N., and Stephanou, E. G.: Formation of atmospheric particles from organic acids produced by forests, *Nature*, 395, 683–686, 1998.
- Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R., Weingartner, E., Frnakevich, V., Zenobi, R., and Baltensberger, U.: Identification of polymers as major components of atmospheric organic aerosols, *Science*, 303, 1659–1662, 2004.
- 25 Kashchiev, D.: On the relation between nucleation work, nucleus size, and nucleation rate, *J. Chem. Phys.*, 76(10), 5098–5102, 1982.
- Kerminen, V.-M. and Kulmala, M.: Analytical formulae connecting the “real” and the “apparent” nucleation rate and the nuclei number concentration for atmospheric nucleation events, *J. Aerosol Science*, 33, 609–622, 2002.
- 30

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

- Kulmala, M., Toivonen, A., Mäkelä, J. M., and Laaksonen, A.: Analysis of the growth of nucleation mode particles observed in Boreal forest, *Tellus*, 50B, 449–462, 1998.
- Kulmala, M., Pirjola, L., and Mäkelä, J. M.: Stable sulphate clusters as a source of new atmospheric particles, *Nature*, 404, 66–69, 2000.
- 5 Kulmala, M., Lauri, A., Vehkamäki, H., Laaksonen, A., Petersen, D., Wagner, P. E.: Strange Predictions by Binary Heterogeneous Nucleation Theory Compared with a Quantitative Experiment, *J. Phys. Chem. B*, 105, 11 800–11 808, 2001.
- Kulmala, M., Hämeri, K., Aalto, P. P., Mäkelä, J. M., Pirjola, L., Nilsson, E. D., Buzorius, G., Rannik, Ü, Dal Maso, M., Seidl, W., Hoffmann, T., Janson, R., Hansson, H.-C., Viisanen, Y.,
10 Laaksonen, A., and O'Dowd, C. D.: Overview of the international project on Biogenic aerosol formation in the boreal forest (BIOFOR), *Tellus*, 53B, 324–343, 2001a.
- Kulmala, M., Dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Miiikkulainen, P., Hämeri, K., and O'Dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus*, 53B, 479–490, 2001b.
- 15 Kulmala, M.: How Particles Nucleate and Grow, *Science*, 302, 1000–1001, 2003.
- Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso, M., Lauri, A., Kerminen, V.-M., Birmili, W., and McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: A review of observations, *J. Aerosol Sci.*, 35, 143–176, 2004a.
- Kulmala, M., Laakso, L., Lehtinen, K. E. J., Riipinen, I., Dal Maso, M., Anttila, T., Kerminen, V.-M., Horrak, U., Vana, M., and Tammet, H.: Initial steps of aerosol growth, *Atmos. Chem. Phys.*, 4, 2553–2560, 2004b,
20 [SRef-ID: 1680-7324/acp/2004-4-2553](#).
- Kulmala, M., Kerminen, V.-M., Anttila, T., Laaksonen, A., and O'Dowd, C. D.: Organic aerosol formation via sulphate cluster activation, *J. Geophys. Res.*, 109(D4), 4205, doi:10.1029/2003JD003961, 2004c.
- 25 Kulmala, M., Lehtinen, K. E. J., Laakso, L., Mordas, G., and Hämeri, K.: On the existence of neutral atmospheric clusters, *Boreal Env. Res.*, 10, 79–87, 2005.
- Laakso, L., Anttila, T., Lehtinen, K. E. J., Aalto, P. P., Kulmala, M., Hörrak, U., Paatero, J., Hanke, M., and Arnold, F.: Kinetic nucleation and ions in boreal particle formation events,
30 *Atmos. Chem. Phys.*, 4, 2353–2366, 2004b,
[SRef-ID: 1680-7324/acp/2004-4-2353](#).
- Laaksonen, A.: Application of nucleation theories to atmospheric aerosol formation, in: *Nucleation and Atmospheric Aerosols 2000*, edited by: Hale, B. N. and Kulmala, M., AIP Confer-

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

- ence Proceedings 534, American Institute of Physics, Melville, NY, pp. 711–723, 2000.
- Lazaridis, M., Kulmala, M., and Gorbunov, B. Z.: Binary heterogeneous nucleation at a non-uniform surface, *J. Aerosol Sci.*, 23, 457–466, 1992.
- Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715–737, 2005,
[SRef-ID: 1680-7324/acp/2005-5-715](#).
- MacDowell, L. G.: Formal study of nucleation as described by fluctuation theory, *J. Chem. Phys.*, 119, 453–463, 2003.
- Mäkelä, J. M., Aalto, P., Jokinen, V., Pohja, T., Nissinen, A., Palmroth, S., Markkanen, T.,
 10 Seitsonen, K., Lihavainen, H., and Kulmala, M.: Observations of ultrafine aerosol particle formation and growth in boreal forest, *Geophys. Res. Lett.*, 24, 1219–1222, 1997.
- Määttänen, A., Vehkamäki, H., Lauri, A., Merikallio, S., Kauhanen, J., Savijärvi, H., and Kulmala, M.: Nucleation studies in the Martian atmosphere, *J. Geophys. Res.*, 110, doi:10.1029/2004JE002308, 2005.
- 15 Mönkkönen, P., Koponen, I. K., Lehtinen, K. E. J., Hämeri, K., Uma, R., and Kulmala, M.: Measurements in a highly polluted Asian mega city: Observations of aerosol aerosol number size distributions, modal parameters and nucleation events, *Atmos. Chem. Phys.*, 5, 57–66, 2005,
[SRef-ID: 1680-7324/acp/2005-5-57](#).
- 20 O'Dowd, C., McFiggins, G., Creasey, D. J., Pirjola, L., Hoell, C., Smith, M. H., Allan, B. J., Plane, J. M. C., Heard, D. E., Lee, J. D., Pilling, M. J., and Kulmala, M.: On the photochemical production of new particles in the coastal boundary layer, *Geophys. Res. Lett.*, 26, 1707–1710, 1999.
- Oxtoby, D. W. and Kashchiev, D.: A general relation between the nucleation work and the size
 25 of the nucleus in multicomponent nucleation, *J. Chem. Phys.*, 100(10), 7665–7671, 1994.
- Pirjola, L., Kulmala, M., Wilck, M., Bischoff, A., Stratmann, F., and Otto, E.: Effects of aerosol dynamics on the formation of sulphuric acid aerosols and cloud condensation nuclei, *J. Aerosol Sci.*, 30, 1079–1094, 1999.
- Vehkamäki, H., Dal Maso, M., Hussein, T., Flanagan, R., Hyvärinen, A., Lauros, J., Merikanto, J., Mönkkönen, P., Pihlatie, M., Salminen, K., Sogacheva, L., Thum, T., Ruuskanen, T.,
 30 Keronen, P., Aalto, P., Hari, P., Lehtinen, K., and Kulmala, M.: Atmospheric particle formation events at Värriö measurement station in Finnish Lapland 1998–2002, *Atmos. Chem. Phys.*, 4, 2015–2023, 2004,

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

- Vehkamäki, H., Napari, I., Kulmala, M., and Noppel, M.: Stable ammonium bisulphate clusters in the atmosphere, *Phys. Rev. Lett.*, 93(14), art. no. 148501, 2004.
- Vehkamäki, H., Lauri, A., Määttänen, A., and Kulmala, M.: Nucleation theorems for heterogeneous nucleation, *Proceedings of European Nucleation Conference 2005*, 28 August–2 September, Ghent, Belgium, p. 34, 2005.
- Viisanen, Y., Strey, R., and Reiss, H.: Homogeneous nucleation rates for water, *J. Chem. Phys.*, 99(6), 4680–4692, 1993.
- Viisanen, Y., Kulmala, M., and Laaksonen, A.: Experiments on gas-liquid nucleation of sulfuric acid and water, *J. Chem. Phys.*, 107, 920–926, 1997.
- Väkevä, M., Hämeri, K., Puhakka, T., Nilsson, E. D., Hohti, H., and Mäkelä, J. M.: Effects of meteorological processes on aerosol particle size distribution in an urban background area, *J. Geophys. Res.*, 105, 9807–9821, 2000.
- Wagner, P. E., Kaller, D., Vrtala, A., Lauri, A., Kulmala, M., and Laaksonen, A.: Nucleation probability in binary heterogeneous nucleation of water-n-propanol vapor mixtures on insoluble and soluble nanoparticles, *Phys. Rev. E*, 67(2), art. no. 021605, Part 1, 2003.
- Weber, R. J., McMurry, P. H., Eisele, F. L., and Tanner, J.: Measurement of expected nucleation precursor species and 3-500-nm diameter particles at Mauna Loa observatory, Hawaii, *J. Atmos. Sci.*, 52, 2242–2257, 1995.
- Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J., and Jefferson, A.: Measured atmospheric new particle formation rates: Implications for nucleation mechanisms. *Chemical Engineering Communications*, 151, 53-64, 1996.
- Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J., and Jefferson, A.: Measurements of new particle formation and ultrafine particle growth rates at a clean continental site, *J. Geophys. Res.*, 102, 4375–4385, 1997.

Formation rate of 3 nm particles

M. Kulmala et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Formation rate of
3 nm particles**

M. Kulmala et al.




Fig. 1. Dependence of apparent nucleation rate on sulphuric acid concentration. J_1 is the particle formation rate at 1 nm, J_3 the apparent particle formation rate at 3 nm and J_3^* the apparent particle formation rate at 3 nm if sulphuric acid does not participate in growth.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

**Formation rate of
3 nm particles**

M. Kulmala et al.




Fig. 2. Small particle concentration (3–6 nm) and scaled and time shifted (90 min ahead) sulphuric acid concentration measured in Hyytiälä in day 84 of the year 2003.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

**Formation rate of
3 nm particles**

M. Kulmala et al.




Fig. 3. Comparison of formation rates of 3 nm particles predicted by different theories and using observed number concentration. Hyytiälä Day 84, 2003.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU